

US005712142A

# United States Patent [19]

Adney et al.

[11] Patent Number: 5,712,142

[45] Date of Patent: Jan. 27, 1998

[54] METHOD FOR INCREASING  
THERMOSTABILITY IN CELLULOSE  
ENZYMES

[75] Inventors: William S. Adney, Golden; Steven R. Thomas, Denver; John O. Baker, Golden; Michael E. Himmel, Littleton; Yat-Chen Chou, Wheat Ridge, all of Colo.

[73] Assignee: Midwest Research Institute, Kansas City, Mo.

[21] Appl. No.: 604,913

[22] Filed: Feb. 22, 1996

## Related U.S. Application Data

[63] Continuation-in-part of Ser. No. 276,213, Jul. 15, 1994, Pat. No. 5,536,655, which is a continuation-in-part of Ser. No. 125,115, Sep. 21, 1993, Pat. No. 5,366,884, which is a continuation-in-part of Ser. No. 826,089, Jan. 27, 1992, Pat. No. 5,275,944, which is a continuation-in-part of Ser. No. 412,434, Sep. 26, 1989, Pat. No. 5,110,735.

[51] Int. Cl.<sup>6</sup> ..... C12N 9/42; C12N 1/20; C12P 21/06; C17H 21/04

[52] U.S. Cl. .... 435/209; 435/69.1; 435/252.3; 435/252.33; 435/320.1; 536/23.1; 536/23.2; 536/23.7

[58] Field of Search ..... 435/209, 69.1, 435/252.3, 252.31, 252.33, 253.5, 254.21, 320.1; 536/22.1, 23.1, 23.2, 23.7

[56] References Cited

## U.S. PATENT DOCUMENTS

5,110,735 5/1992 Tucker et al. .... 435/209

5,275,944 1/1994 Himmel et al. .... 435/209  
5,366,884 11/1994 Adney et al. .... 435/209  
5,432,075 7/1995 Himmel et al. .... 435/209  
5,536,655 7/1996 Thomas et al. .... 435/209

## OTHER PUBLICATIONS

Mohagheghi et al., *Int. J. System. Bacteriol.*, 36:435-443 (1986).

Lejeune et al., *Biosynthesis and Biodegradation of Cellulose*, C. Haigler and P.J. Weimer, Ed., Marcel-Dekker, NY 1991, pp. 623-672.

Primary Examiner—Rebecca E. Prouty

Assistant Examiner—Tekchand Saidha

Attorney, Agent, or Firm—Edna M. O'Connor; Ken Richardson; Ruth Eure

[57] ABSTRACT

The gene encoding *Acidothermus cellulolyticus* E1 endoglucanase is cloned and expressed in *Pichia pastoris*. A new modified E1 endoglucanase enzyme comprising the catalytic domain of the full size E1 enzyme demonstrates enhanced thermostability and is produced by two methods. The first method of producing the new modified E1 is proteolytic cleavage to remove the cellulose binding domain and linker peptide of the full size E1. The second method of producing the new modified E1 is genetic truncation of the gene encoding the full size E1 so that the catalytic domain is expressed in the expression product.

13 Claims, 8 Drawing Sheets




Fig. 1




Fig. 2




Fig. 3




Fig. 4

AGGGYWHTSG REILDANNVP VRIAGINWFG FETCNVYVHG LWSRDYRSM	50
DQIKSLGYNT IRLPYSDDIL KPGTMPNSIN FYQMNQDLQG LTSLQVMDKI	100
VAYAGQIGLR IILDRHRPDC SGQSALWYTS SVSEATWISD LQALAORYKG	150
NPTVVGFDLH NEPHDPACWG CGDPSIDWRL AAERAGNAVL SVNPNLLIFV	200
EGVQSYNGDS YWWGGNLQGA GQYPVVLNVP NRLVYSAHDY ATSVYPQTF	250
SDPTFPNNMP GIWNKNWGYL FNQNIAPVWL GEFGTTLQST TDQTWLKT	300
QYLRPTAQYG ADSFQWTFWS WNPDSGDTGG ILKDDWQTV	350
KSSIFDPV	358

**Fig. 5**

GGATCCACGT TGTACAAGGT CACCTGTCCG TCGTTCTGGT AGAGCGGCGG GATGGTCACC	60
CGCACGATCT CTCCTTTGTT GATGTGACG GTCACGTGGT TACGGTTTGC CTCGGCCGCG	120
ATTTTCGCGC TCGGGCTTGC TCCGGCTGTC GGGTTCGGTT TGGCGTGGTG TGCGGAGCAC	180
GCCGAGGCGA TCCCAATGAG GGCAAGGGCA AGAGCGGAGC CGATGGCACG TCGGGTGGCC	240
GATGGGGTAC GCCGATGGGG CGTGGCGTCC CCGCCGCGGA CAGAACCGBA TGCGGAATAG	300
GTCACGGTGC GACATGTTGC CGTACCGCGG ACCCGGATGA CAAGGGTGGG TGCGCGGGTC	360
GCCTGTGAGC TGCCGGCTGG CGTCTGGATC ATGGGAACGA TCCCACCATT CCCCACCAATC	420
GACGCGATCG GGAGCAGGGC GCGCGGAGCC GGACCGTGTG GTCGAGCCGG ACGATTGCGC	480
CATACGGTGC TGCAATGCC AGCGCCATGT TGTCAATCCG CCAAATGCAG CAATGCACAC	540
ATGGACAGGG ATTGTGACTC TGAGTAATGA TTGGATTGCC TTCTTGCCGC CTACGCGTTA	600
CGCAGAGTAG GCGACTGTAT GCGGTAGGTT GCGCTCCAG CCGTGGGCTG GACATGCCTG	660
CTGCGAACTC TTGACACGTC TGGTTGAACG CGCAATACTC CCAACACCGA TGGGATCGTT	720
CCCATAAGTT TCCGTCTCAC AACAGAATCG GTGCGCCCTC ATGATCAACG TGAAGGAGT	780
ACGGGGGAGA ACAGACGGGG GAGAAACCAA CGGGGGATTG GCGGTGCCGC GCGCATTGCG	840
GCGAGTGCTT GGCTCGCGGG TGATGCTGCG GGTGCGGCGT GTCGTCGCGG TGCTGGCATT	900
GGTTGCCGCA CTCGCCAACC TAGCCGTGCC GCGGCCGGCT CCGCCGCGG GCGGCGGCTA	960
TTGGCACACG AGCGGCCGGG AGATCCTGGA CGCGAACAAC GTGCCGGTAC GGATCGCCGG	1020
CATCAACTGG TTTGGGTTCC AAACCTGCAA TTACGTCGTG CACGGTCTCT GGTCACGCGA	1080
CTACCGCAGC ATGCTCGACC AGATAAAGTC GCTCGGCTAC AACACAATCC GGCTGCCGTA	1140
CTCTGACGAC ATTCTCAAGC CGGGCACCAT GCCGAACAGC ATCAATTTT ACCAGATGAA	1200
TCAGGACCTG CAGGGTCTGA CGTCCTTGCA GGTTCATGGAC AAAATCGTCG CGTACGCCGG	1260
TCAGATCGGC CTGCGCATCA TTCTTGACCG CCACCGACCG GATTGCAGCG GGCAGTCGGC	1320
GCTGTGGTAC ACGAGCAGCG TCTCGGAGGC TACGTGGATT TCCGACCTGC AAGCGCTGGC	1380
GCAGCGCTAC AAGGGAAACC CGACGGTCGT CGGCTTTGAC TTGCACAACG AGCCGCATGA	1440
CCCGGCCTGC TGGGGCTGCG GCGATCCGAG CATCGACTGG CGATTGGCCG CCGAGCGGGC	1500
CGGAAACGCC GTGCTCTCGG TGAATCCGAA CCTGCTCATT TTCGTCGAAG GTGTGCAGAG	1560
CTACAACGGA GACTCCTACT GGTGGGGCGG CAACCTGCAA GGAGCCGGCC AGTACCCGGT	1620
CGTGCTGAAC GTGCCGAACC GCCTGGTGTA CTCGGCGCAC GACTACGCGA CGAGCGTCTA	1680
CCCGCAGACG TGGTTCAGCG ATCCGACCTT CCCCACAAC ATGCCCGGCA TCTGGAACAA	1740
GAAGTGGGGA TACCTCTTCA ATCAGAACAT TGCACCGGTA TGGCTGGGCG AATTTCGGTAC	1800
GACACTGCAA TCCACGACCG ACCAGACGTG GCTGAAGACG CTCGTCCAGT ACCTACGGCC	1860
GACCGCGCAA TACGGTGGG ACAGCTTCCA GTGGACCTTC TGGTCTGGA ACCCCGATTC	1920
CGGCGACACA GGAGGAATTC TCAAGGATGA CTGGCAGACG GTCGACACAG TAAAGACGG	1980
CTATCTCGCG CCGATCAAGT CGTCGATTTT CGATCCTGTC <u>TAATGATCTG CATCGCCTAG</u>	2040
<u>CAGTCAACCG</u> TCCCCGTCGG TGTCGCGGTC TCCGTCGCGG AGCCCGTCGG CGAGTCGGAC	2100
GCCGACGCCT ACTCCGACGC CGACAGCCAG CCCGACGCCA ACGCTGACCC CTACTGCTAC	2160
GCCCACGCC ACGGCAAGCC CGACGCCGTC ACCGACGGCA GCCTCCGGAG CCCGCTGCAC	2220
CGCGAGTTAC CAGGTCAACA GCGATTGGGG CAATGGCTTC ACGGTAACGG TGGCCGTGAC	2280
AAATTCCGGA TCC	2293

Fig. 6




Fig. 7






Fig. 8

## METHOD FOR INCREASING THERMOSTABILITY IN CELLULOSE ENNZYMES

This application is a continuation-in-part of Ser. No. 08/276,213, filed Jul. 15, 1994, now U.S. Pat. No. 5,536,655, which is a continuation-in-part of Ser. No. 08/125,115, filed Sep. 21, 1993, now U.S. Pat. No. 5,366,884, which is a continuation-in-part of Ser. No. 07/826,089, filed Jan. 27, 1992, now U.S. Pat. No. 5,275,944, which was a continuation-in-part of Ser. No. 412,434, filed Sep. 26, 1989 now U.S. Pat. No. 5,110,735.

The United States Government has rights in this invention under Contract No. DE-AC36-83CH10093 between the United States Department of Energy and the National Renewable Energy Laboratory, a Division of the Midwest Research Institute.

### FIELD OF THE INVENTION

The invention relates to methods for increasing the thermostability of cellulose enzymes.

### BACKGROUND OF THE INVENTION

The development of an economic process for the conversion of low-value biomass to useful products via fermentation requires the optimization of several key steps, including cellulase production and performance. Practical utilization of cellulose by hydrolysis with cellulase to produce glucose requires large amounts of cellulase to fully depolymerize cellulose. For example, about one kilogram cellulase preparation may be used for every fifty kilograms of cellulose. Economical production of cellulose is also compounded by the relatively slow growth rates of cellulase-producing fungi and the long times required for cellulase induction. Therefore, improvements in or alternative cellulase production systems capable of greater productivities of cellulase activity than may be possible from currently available systems would significantly reduce the cost of cellulose hydrolysis and make the large-scale bioconversion of cellulose biomass more economical.

Highly thermostable cellulose enzymes are secreted by the cellulolytic thermophile *Acidothermus cellulolyticus* gen. nov., sp. nov. These are discussed in U.S. Pat. Nos. 5,110,735, 5,275,944, 5,366,884, and 5,432,075. All four of these patents are included herein in their entirety by this reference thereto. This bacterium was originally isolated from decaying wood in an acidic, thermal pool at Yellowstone National Park and deposited with the American Type Culture Collection (ATCC) under collection number 43068 (Mohagheghi et al. 1986. *Int. J. System. Bacteriol.* 36:435-443).

The cellulase complex produced by this organism is known to contain several different cellulase enzymes with maximal activities at temperatures of 75° C. to 83° C. These cellulases are resistant to inhibition from cellobiose, an end product of the reactions catalyzed by cellulase. Also, the cellulases from *Acidothermus cellulolyticus* are active over a broad pH range centered about pH 6. A high molecular weight cellulase isolated from growth broths of *Acidothermus cellulolyticus* was found to have a molecular weight of approximately 156,600 to 203,400 daltons by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). This enzyme is described in U.S. Pat. No. 5,110,735.

A novel cellulase enzyme, known as the E1 endoglucanase, also secreted by *Acidothermus cellulolyticus* into the growth medium, is described in detail in U.S. Pat.

No. 5,275,944. In its native form, this endoglucanase demonstrates a temperature optimum of 83° C. and a specific activity of 40  $\mu$ mole glucose release from carboxymethylcellulose/min/mg protein. This E1 endoglucanase was further identified as having an isoelectric pH of 6.7. It is this E1 endoglucanase which has been modified and made the subject of this patent application. The E1 endoglucanase is a multidomain cellulase having a catalytic domain and a cellulose binding domain connected to the catalytic domain by a linker peptide.

### SUMMARY OF THE INVENTION

The E1 endoglucanase described above, has been modified to increase its thermostability. The present modification has increased the thermostability of this enzyme by effectively doubling the length of time which this enzyme demonstrates half-maximal activity at elevated temperatures, as well as increasing the temperature at which maximal rates of catalysis are observed. The modification comprises eliminating the cellulose binding domain and linker peptide of the enzyme from the catalytic domain. It is the catalytic domain containing the catalytically active portion of the molecule, which remains after elimination of the cellulose binding domain and linker peptide, which demonstrates these improved thermal properties. This modification has been accomplished by two methods. The first method for eliminating the cellulose binding domain and linker peptide is by subjecting the entire molecule to proteolytic cleavage, which removes the cellulose binding domain and linker peptide from the catalytic domain. The second method for removing the cellulose binding domain and linker peptide from full size E1 involves modification of the gene which encodes full size E1 so that the cellulose binding domain and linker peptide are not present in the expression product. The E1 enzyme of the present invention which demonstrates enhanced thermostability by elimination of the cellulose binding domain and linker peptide is referred to as "modified", "truncated", "catalytic domain", or "E1 CAT". Also, E1 CAT produced by proteolytic cleavage may be referred to as pE1 CAT and E1 CAT produced by genetic transformation may be referred to as gel CAT.

In addition to the modified E1 endoglucanase having improved thermostability, this invention teaches the expression of full size E1 in a yeast

It is therefore an object of the present invention to transform and express the full size E1 endoglucanase gene in a yeast under the same and/or a different gene regulatory system.

It is an object of the present invention to modify the gene encoding the E1 endoglucanase from *Acidothermus cellulolyticus* to enhance its thermostability by eliminating expression of the cellulose binding domain and linker peptide in the gene product.

It is an object of the present invention to provide the DNA sequence which encodes the modified form of the E1 endoglucanase from *Acidothermus cellulolyticus*.

It is another object of the present invention to provide the amino acid sequence of the modified form of the E1 endoglucanase from *Acidothermus cellulolyticus*.

It is another object of the present invention to provide a method for proteolytic cleavage of the cellulose binding domain and linker peptide from the E1 endoglucanase.

It is a further object of the present invention to prepare modified E1 endoglucanases which have different properties from the natural enzyme.

The present invention describes the gene for and the nucleotide sequence of the segment of *Acidothermus cellu-*

*lotyticus* DNA encoding the catalytic domain of the E1 endoglucanase gene. This 2293 base fragment of DNA is unique in nature and discretely defined. The natural gene contains a promoter, a ribosome binding site, a signal peptide, an open reading frame, a termination codon and a putative transcriptional terminator. The modified gene contains a promoter, a ribosome binding site, a signal peptide and one or more termination codons inserted at the C terminus of the catalytic domain.

The cloned gene may also be expressed in other microorganisms under its natural promoter or another promoter recognized by the host microorganism. The cloned gene may be expressed in mammalian systems, higher plants or vital vectors. Alternatively, additional copies of the gene may be introduced into *Acidothermus cellulolyticus* or other heterologous host organism to enhance expression of the enzyme. Additionally, DNA encoding one or more domains of the *Acidothermus cellulolyticus* E1 endoglucanase may be ligated to domains in other compatible endoglucanases to make a recombinant DNA capable of expressing a hybrid endoglucanase enzyme having beneficial properties from both endoglucanases.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a plasmid map of pPIC9-E1.

FIG. 2 compares the functional half-life of the full size E1 enzyme to the functional half-life of the catalytic domain of the E1 enzyme at 80° C.

FIG. 3 compares the temperature optimum of the full size E1 enzyme to the temperature optimum of the catalytic domain of the E1 enzyme.

FIG. 4 is a chromatogram showing purified constituent domains of full size E1 by size exclusion chromatography.

FIG. 5 shows the amino acid translation of the coding sequence of E1 CAT.

FIG. 6 shows the 2293 base pair nucleotide sequence of the region of *Acidothermus cellulolyticus* genomic DNA which contains the modified E1 endoglucanase gene which expresses only the catalytic domain of the enzyme, without the linker peptide or cellulose binding domain.

FIG. 7 is a plasmid map of pYCC101.

FIG. 8 is a DSC thermogram comparing the denaturation endotherm peak at 78° C. for the full size E1 and E1 CAT showing no peak below at least 88° C.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

According to the present invention the entire E1 coding sequence for *Acidothermus cellulolyticus* E1 endoglucanase is cloned and expressed in a different microbial host than is described in the parent to this application, Ser. No. 08/276,213, filed Jul. 15, 1994, which is incorporated herein in its entirety by this reference thereto. The enzyme described in Ser. No. 08/276,213 is a  $\beta$ -1,4 endoglucanase which can hydrolyze cellulose or carboxymethylcellulose and is hereafter referred to as E1 endoglucanase or full size E1. The result is a vastly improved rate of E1 enzyme production over native or genetically engineered bacterial systems, thereby lowering the cost of cellulase. The instant application teaches expression of full size E1 in a yeast, namely *Pichia pastoris*.

#### Expression of Full Size E1 and Modified E1

For expressing the E1 endoglucanase gene, either the full size E1 gene or the modified E1 gene, one may use a variety

of hosts including most bacteria, yeast, fungi, algae, viruses, plants and animals. Organisms which naturally produce cellulase enzymes are preferred host cells along with easy to grow host cells and host cells known to express and secrete heterologous genes in large quantities.

If the host cell is a bacterium, generally a bacterial promoter and regulatory system will be used. For a typical bacterium such as *E. coli*, representative examples of well known promoters include, for example, trc, lac, tac, trp, bacteriophage lambda P<sub>L</sub>, T7 RNA polymerase promoter, etc. When the expression system is yeast, examples of well known promoters include, but are not limited to GAL 1/GAL 10, alcohol dehydrogenase (ADH), alcohol oxidase (AOX), his3, cyc1, etc. For eukaryotic hosts, enhancers such as the yeast Ty enhancer, may be used.

Alternatively, if one wished for the full size or modified E1 endoglucanase gene to be expressed at only a particular time, such as after the culture or host organism has reached maturity, an externally regulated, environmentally-responsive promoter is particularly useful. Examples include those based upon the nutritional or chemical composition of the medium (e.g. methanol, lac, trp, his), temperature regulation (e.g. temperature sensitive regulatory elements), heat shock promoters (e.g. HSP80A, U.S. Pat. No. 5,187,267), stress response (e.g. plant EF1A promoter, U.S. Pat. No. 5,177,011) and chemically inducible promoters (e.g. tetracycline inducible promoter or salicylate inducible promoter U.S. Pat. No. 5,057,422).

Other suitable hosts for expressing full size or modified E1 endoglucanase include members of the genera: *Trichoderma*, *Fusarium*, *Penicillium*, *Bacillus*, *Xanthomonas*, *Streptomyces*, *Aspergillus* and *Pichia*, for example. Some of these microorganisms also serve as sources of endoglucanase genes for the formation of mixed domain genes for the production of hybrid enzymes.

Expression of the full size E1 endoglucanase gene has been demonstrated in *E. coli*, *Pichia pastoris* and in *Streptomyces lividans*.

Expressing full size or modified E1 endoglucanase in *E. coli* may be performed under control of a T7 bacteriophage promoter or other promoter recognizable by *E. coli*. Expression of full size E1 in *E. coli* has been enhanced considerably relative to the native gene with the constructs of the present invention. Expression of the full size E1 endoglucanase coding sequence in *S. lividans* has been achieved twice more with several different constructs employing two different promoters. These are the tipA promoter (thiostrepton-inducible) and the ST I-II promoter isolated from a trypsin inhibitor gene from *Streptomyces longisporus*. Expression levels of active, full size, secreted E1 endoglucanase up to 20 mg/L have been achieved with the ST I-II promoter.

Expression of the full size or modified E1 endoglucanase coding sequence in the filamentous fungi, *Aspergillus niger*, *A. awamori*, *A. oryzae*, *A. terreus* and/or *A. nidulans* and *Trichoderma reesei* is achievable using various promoters derived from *Aspergillus* or *Trichoderma*. These promoters include, but are not limited to, G3PDH (glyceraldehyde-3-phosphate dehydrogenase), glucoamylase,  $\beta$ -tubulin and IPNS (isopenicillin N synthase) from *Aspergillus* and CBH I (cellobiohydrolase I), alcohol dehydrogenase, triosephosphate isomerase and  $\alpha$ -amylase from *T. reesei*.

The *Acidothermus cellulolyticus* full size E1 endoglucanase gene was cloned, as described in Ser. No. 08/276,213, and expressed in *Pichia pastoris* using the AOX1 promoter and 3' sequences. This expression is taught in Example 1.

#### EXAMPLE 1

##### Expression of the Entire Full Size E1 Gene in *Pichia pastoris*

Expression of active E1 endoglucanase in the range of 0.75–1.5 g/L has been accomplished in the yeast, *Pichia*

Plasmid 4-5, a pGEM-7 (Promega Corp.) derivative carrying a 3.7 kb genomic fragment of *Acidothermus cellulyticus* DNA and harboring the entire E1 gene, was used as a template in PCR reactions to amplify the E1 coding sequence for subsequent cloning into the *Pichia* secretion vector, pPIC9. The primer annealing to the non-coding strand of the template, "E1-f", is a 30-mer with 18 bases of homology to the template, beginning at the mature N-terminus of the E1 polypeptide. The segment of E1-f which is not homologous to the template molecules incorporates 4 codons which encode the C-terminal 4 amino acids of the  $\alpha$ F signal peptide which is present in pPIC9 and also includes an XhoI site at the 5' end for use in subsequent cloning into pPIC9. The primer annealing to the coding strand and priming synthesis of the non-coding strand, "E1-r", is a 24-mer with 18 bases of homology to the template. The sequence of E1-r corresponds to the last 5 codons of the E1 coding sequence and the stop codon. The non-homologous 5' tail of E1-r adds an AvrII restriction site for use in subsequent cloning into pPIC9.

The media and intracellular contents of cells from cultures grown in the presence of methanol for two days were screened by Western blot analysis using a monoclonal antibody specific for the E1 endoglucanase. All *P. pastoris* clones containing the foreign DNA were shown to express

This enhancement is brought about by the cleavage of the cellulose binding domain from the catalytic domain of the full size E1 enzyme. The cleavage of the cellulose binding

The recombinant full size enzyme was purified with essentially a three step purification process consisting of hydrophobic interaction chromatography (HIC) followed by anion exchange and finally, size exclusion chromatography. The HIC step employed a column packed with 250 mL of Pharmacia Fast Flow Phenyl Sepharose. This was followed

by anion-exchange chromatography using a 6 mL Pharmacia Resource Q anion exchange column.

For the hydrophobic interaction step, ammonium sulfate was added to the concentrated culture supernatant of one 10 L fermentation to a final concentration 0.5M. A total volume of 300 mL of this concentrate was loaded onto the phenyl sepharose column and washed extensively with 20 mM Tris, 0.5M (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> pH 8.0. The column was developed with a linear decreasing gradient (0.5M-zero) of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>. Recombinant full size E1 eluted at zero salt concentration. Fractions containing E1 activity were identified using 4-methylumbelliferyl  $\beta$ -D-cellobioside (MUG) assays. Active fractions were combined, concentrated and diafiltered against 20 mM Tris pH 8.0, after which they were loaded directly onto the anion-exchange column and eluted with a increasing NaCl gradient (0-300 mM). A final buffer exchange and purification step was done using size exclusion chromatography with a 2.6 cm $\times$ 10 cm Pharmacia Superdex 200 column and using one of two buffer systems; either 20 mM acetate, 100 mM NaCl, pH 5.0 buffer or 50 mM ammonium acetate, pH 6.2 buffer depending upon the eventual use of the enzyme.

The E1 catalytic domain was produced by proteolytic cleavage with papain. The proteolytic digestions were done in 50 mM ammonium acetate, pH 6.2 buffer. Molar ratios of 72 kDa MW E1/23 kDa MW papain cleaved E1 of 6/1 were used. Papain digestions were incubated at 28° C. for 24 h. The catalytic domain was separated from full size recombinant E1, E1-CBD (cellulose binding domain), linker peptides and papain by SEC using a 2.6 cm $\times$ 10 cm Pharmacia Superdex 200 column.

A chromatogram demonstrating the purification of modified catalytic domain from papain cleaved *S. lividans* recombinant E1 by size exclusion chromatography is shown in FIG. 4. Peak A is unmodified full size E1 enzyme, peak B is the catalytic domain, and peak C is the cellulose binding domain of the cleaved product. Also, peak B reacts with a monoclonal antibody (MAB) specific for full size E1, thereby confirming that this fragment contains epitopes for which this MAB shows specificity.

In order to confirm the identity of the peptide isolated from the papain digestion, the peak B peptide was subjected to analytical ultracentrifugation using a Beckman Optima XLA centrifuge. Sedimentation equilibrium analysis yielded an estimated molecular weight of 41,600 daltons for this peptide. SDS-PAGE analysis rendered a molecular weight estimate of 42,000 daltons. These values compare well with the calculated molecular weight predicted from amino acid sequence for the catalytic domain (i.e., 40,192 da.) Due to the limits of detection of the techniques utilized herein, these molecular weight values are well within the range of experimental error and therefore the molecular weight of the E1 CAT is in the range of about 40,000 to 42,000 Daltons.

Comparison of the thermal stability at 80° C. of pE1 CAT and full size E1 endoglucanase, both produced from a *Streptomyces* host, can be seen in FIG. 2. The enzyme was

activity using 1 mg/mL p-nitrophenyl- $\beta$ -D-cellobioside substrate in 20 mM acetate, 100 mM NaCl, pH 5.0, at 65° C. for 30 minutes. Activity values are expressed as a percentage of the activity detected at time zero (e.g., 100%).

Comparison of the temperature optima of modified E1 and full size E1 was performed using 1 mg/mL p-nitrophenyl-g-D cellobioside substrate in 20 mM acetate, 100 mM NaCl, pH 5.0, at various temperatures with a 30 minute incubation time. See FIG. 3. The temperature optimum of the E1 CAT produced by papain cleavage is increased by 10° C. relative to the full size E1. Activity values in FIG. 3 are expressed as a percentage of the maximum activity detected.

It is further contemplated that one may include more than one catalytic domain in the hybrid enzyme. This may allow for a further increase in specific activity. Also, a catalytic domain containing cellulase activity other than endoglucanase activity may be included as well to reduce the number of cellulase enzymes one needs to add to a cellulosic substrate for polymer degradation.

### EXAMPLE 3

#### Production of E1 CAT by Genetic Truncation of the Full Size E1 Coding Sequence

The amino acid sequence of E1 CAT is shown in FIG. 5. The entire amino acid sequence of pE1 CAT, including the C-terminus, is confirmed by the x-ray crystal structure derived from this molecule. An alternative to production of E1 CAT by papain cleavage of full size E1 is taught A molecular genetic approach was also developed to produce E1 CAT.

A strategy was desired to generate a genetically truncated E1 gene which would produce E1 CAT without requiring any downstream processing to achieve E1 CAT from a precursor molecule, as for papain cleavage. One way to accomplish this is to introduce a translational stop codon at or near the C-terminal residue of the catalytic domain.

The 2.3 kb Bam HI fragment containing most of the E1 gene was subcloned into pAlter-1 in preparation for site-directed mutagenesis (pYCC100). The nucleotide sequence for this fragment is shown in FIG. 6. A mutagenic 36-mer oligonucleotide (underlined in FIG. 6) was synthesized (5'-ATTTTCGATC CTGTCTAATG ATCTGCATCG CCTAGC-3')(SEQ ID NO:5). Using the Altered Sites® II kit (Promega Corp.) two consecutive codons immediately downstream of the C-terminal residue of E1-CAT (as determined by x-ray crystallography) were changed to different stop codons (TAA, TGA). The six mutagenic nucleotides are double-underlined in FIG. 6. The DNA sequence of this clone (pYCC101) has been confirmed by dideoxy DNA sequencing in the region of the site-directed mutations using the T7 Sequenase kit supplied by US Biochemical. (Cleveland, Ohio). A plasmid map of pYCC101 is shown in FIG. 7.

Native AA seq	S S I F D P V G A S A S P S S Q (SEQ ID NO: 6)
Native DNA seq	TCGTCGATTTTCGATCCTGTGCGCGCTGTCATCGCCTACGAGTCAA (SEQ ID NO: 7)
Mutagenic oligo	ATTTTCGATCCTGTCTAATGATCTGCATCGCCTAGC (SEQ ID NO: 8)
Mutated DNA seq	TCGTCGATTTTCGATCCTGTCTAATGATCTGCATCGCCTAGCAGTCAA (SEQ ID NO: 9)
Mutated AA seq	S S I F D P V . . S A S P S S Q (SEQ ID NO: 10)

incubated at 80° C. in 20 mM acetate, 100 mM NaCl, pH 5.0 buffer with timed aliquots removed, and each assayed for

Mutagenized DNA was transformed into *E. coli* strain ES1301. Transformants were screened for resistance to

ampicillin and sensitivity to tetracycline in order to identify clones carrying the putatively mutagenized E1 gene. Many ampicillin-resistant candidate clones were subsequently screened on plates containing 1 mM 4-methylumbelliferyl- $\beta$ -D-cellobioside (MUC) to verify expression of active E1. Plasmid DNA was prepared from several clones and employed as templates in dideoxy DNA sequencing reactions using the Sequenase® kit (U.S. Biochemical, Cleveland, Ohio) to verify the sequence of E1 DNA in the region of the intended mutation. The mutated sequence was detected in every clone which was sequenced. One of these clones was selected and designated pYCC101. Each of the successfully mutated clones expresses a protein not present in control cells and which migrates at a molecular weight of approximately 42 kDa in SDS-PAGE gels. This 42 kDa protein also reacts with a monoclonal antibody specific for the E1 endoglucanase on Western blots, thus confirming its identity as E1 CAT.

#### EXAMPLE 4

##### Differential Scanning Calorimetry (DSC)

Calorimetric studies of the denaturation of the full size E1 enzyme and the proteolytically cleaved E1 CAT were carried at pH 5.0 in 50 mM sodium acetate, using a Microcal MC-2 differential scanning microcalorimeter over a temperature range of 25°–95° C. and using a scan rate of 20° C./h. For the examples shown in FIG. 2, the protein concentrations were 0.24 mg/mL for the native E1 enzyme and 0.14 mg/mL for E1 CAT.

FIG. 8 shows the DSC thermogram for full size E1 enzyme displaying a prominent denaturation endotherm peak at approximately 78° C. This compares to the thermogram of E1 CAT, which shows no peak below at least 88° C. Since the thermograms are uninterpretable above 88° C. (limitation of the instrumentation), a reasonable conclusion is that the denaturation endotherm for the catalytic domain lies somewhere above 88° C.

#### EXAMPLE 5

##### Method for the Determination of the Functional Half-life of E1 and E1 CAT

The time at which half of the original endoglucanase activity remains following pre-incubation at 80° C. is

referred to as its functional half-life. Enzymes were pre-incubated at 80° C. in 20 mM acetate, 100 mM NaCl, pH 5.0, in concentrations of 13.14  $\mu$ g/mL for the full size enzyme E1 and 19.53  $\mu$ g/mL for E1 CAT. Small aliquots (100  $\mu$ L) from single tubes were removed at various times and assayed for activity by adding the 100  $\mu$ L enzyme aliquot to 700  $\mu$ L 20 mM acetate, 100 mM NaCl, pH 5.0, and 200  $\mu$ L of 5 mg/mL p-nitrophenyl  $\beta$ -D cellobioside. The assay mixture was incubated at 65° C. for 30 minutes and the reaction stopped by adjusting the pH by addition of 2 mL of 1M Na<sub>2</sub>CO<sub>3</sub>. The activity was then measured by determining the concentration of the nitrophenolate anion, released as a result of catalytic activity of the enzyme, by measuring the absorbance at 410 nm of the quenched samples. Results shown in FIG. 2 demonstrate that E1 CAT has a functional half-life at 80° C. which nearly doubles that of full size E1 (16 h vs. 9 h).

Other options for generating genetically truncated E1 contemplated by the present inventors to be part of this invention include PCR of the coding sequence incorporating a non-homologous stop codon into the downstream synthetic primer; or using available restriction sites downstream of the DNA encoding the catalytic domain to delete the DNA sequences encoding the linker peptide and cellulose binding domain. Unless specifically defined otherwise, all technical or scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials have been described.

The foregoing description of the specific embodiments reveal the general nature of the invention so that others can, by applying current knowledge, readily modify and/or adapt for various applications such specific embodiments without departing from the generic concept, and, therefore, such adaptations and modifications should and are intended to be comprehended within the meaning and range of equivalents of the disclosed embodiments. It is to be understood that the phraseology or terminology employed herein is for the purpose of description and not of limitation.

All references mentioned in this application are incorporated by reference.

#### SEQUENCE LISTING

##### ( 1 ) GENERAL INFORMATION:

( i i i ) NUMBER OF SEQUENCES: 12

##### ( 2 ) INFORMATION FOR SEQ ID NO:1:

###### ( i ) SEQUENCE CHARACTERISTICS:

- ( A ) LENGTH: 30 base pairs
- ( B ) TYPE: nucleic acid
- ( C ) STRANDEDNESS: double
- ( D ) TOPOLOGY: linear

( i i ) MOLECULE TYPE: DNA

###### ( i x ) FEATURE:

- ( A ) NAME/KEY: E1-f primer

( x i ) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

-continued

CTC GAG AAA AGA GCG GGC GGC GGC TAT TGG 30

## ( 2 ) INFORMATION FOR SEQ ID NO: 2:

- ( i ) SEQUENCE CHARACTERISTICS:
  - ( A ) LENGTH: 10 amino acids
  - ( B ) TYPE: amino acid
  - ( C ) STRANDEDNESS: single
  - ( D ) TOPOLOGY: linear

( i i ) MOLECULE TYPE: Protein

- ( i x ) FEATURE:
  - ( A ) NAME/KEY: E1-f primer

( x i ) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Leu Glu Lys Arg Ala Gly Gly Gly Tyr Trp  
5 10

## ( 2 ) INFORMATION FOR SEQ ID NO: 3:

- ( i ) SEQUENCE CHARACTERISTICS:
  - ( A ) LENGTH: 24 base pairs
  - ( B ) TYPE: nucleic Acid
  - ( C ) STRANDEDNESS: double
  - ( D ) TOPOLOGY: linear

( i i ) MOLECULE TYPE: DNA

- ( i x ) FEATURE:
  - ( A ) NAME/KEY: E1r

( x i ) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CCT AGG TTA ACT TGC TGC GCA GGC 24

## ( 2 ) INFORMATION FOR SEQ ID NO: 4:

- ( i ) SEQUENCE CHARACTERISTICS:
  - ( A ) LENGTH: 5 amino acids
  - ( B ) TYPE: amino acid
  - ( C ) STRANDEDNESS: single
  - ( D ) TOPOLOGY: linear

( i i ) MOLECULE TYPE: Protein

- ( i x ) FEATURE:
  - ( A ) NAME/KEY: E1r

( x i ) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Ser Ala Ala Cys Ala  
5

## ( 2 ) INFORMATION FOR SEQ ID NO: 5:

- ( i ) SEQUENCE CHARACTERISTICS:
  - ( A ) LENGTH: 36 base pairs
  - ( B ) TYPE: nucleic Acid
  - ( C ) STRANDEDNESS: double
  - ( D ) TOPOLOGY: linear

( i i ) MOLECULE TYPE: DNA

( x i ) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

ATTTTCGATC CTGTCTAATG ATCTGCATCG CCTAGC 36

## ( 2 ) INFORMATION FOR SEQ ID NO: 6:

- ( i ) SEQUENCE CHARACTERISTICS:
  - ( A ) LENGTH: 16 amino acids
  - ( B ) TYPE: amino acid

-continued

( C ) STRANDEDNESS: single  
( D ) TOPOLOGY: linear

( i i ) MOLECULE TYPE: Protein

( x i ) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Ser Ser Ile Phe Asp Pro Val Gly Ala Ser Ala Ser Pro Ser Ser Gln  
5 10 15

( 2 ) INFORMATION FOR SEQ ID NO: 7:

( i ) SEQUENCE CHARACTERISTICS:  
( A ) LENGTH: 48 base pairs  
( B ) TYPE: nucleic acid  
( C ) STRANDEDNESS: double  
( D ) TOPOLOGY: linear

( i i ) MOLECULE TYPE: DNA

( x i ) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

TCGTCGATTT TCGATCCTGT CGGCGCGTCT GCATCGCCTA GCAGTCAA 48

( 2 ) INFORMATION FOR SEQ ID NO: 8:

( i ) SEQUENCE CHARACTERISTICS:  
( A ) LENGTH: 36 base pairs  
( B ) TYPE: nucleic acid  
( C ) STRANDEDNESS: double  
( D ) TOPOLOGY: linear

( i i ) MOLECULE TYPE: DNA

( i x ) FEATURE:

( A ) NAME/KEY: mutagenic oligo

( x i ) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

ATTTTCGATC CTGTCTAATG ATCTGCATCG CCTAGC 36

( 2 ) INFORMATION FOR SEQ ID NO: 9:

( i ) SEQUENCE CHARACTERISTICS:  
( A ) LENGTH: 48 base pairs  
( B ) TYPE: nucleic acid  
( C ) STRANDEDNESS: double  
( D ) TOPOLOGY: linear

( i i ) MOLECULE TYPE: DNA

( i x ) FEATURE:

( A ) NAME/KEY: mutated DNA

( x i ) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

TCGTCGATTT TCGATCCTGT CTAATGATCT GCATCGCCTA GCAGTCAA 48

( 2 ) INFORMATION FOR SEQ ID NO: 10:

( i ) SEQUENCE CHARACTERISTICS:  
( A ) LENGTH: 16 amino acids  
( B ) TYPE: amino acid  
( C ) STRANDEDNESS: single  
( D ) TOPOLOGY: linear

( i i ) MOLECULE TYPE: Protein

( i x ) FEATURE:

( A ) NAME/KEY: Mutated amino acid

( x i ) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Ser Ser Ile Phe Asp Pro Val Xaa Xaa Ser Ala Ser Pro Ser Ser Gln  
5 10 15



-continued

## ( 2 ) INFORMATION FOR SEQ ID NO: 11:

## ( i ) SEQUENCE CHARACTERISTICS:

- ( A ) LENGTH: 358 amino acids
- ( B ) TYPE: amino acid
- ( C ) STRANDEDNESS: single
- ( D ) TOPOLOGY: linear

## ( i i ) MOLECULE TYPE: Protein

## ( i x ) FEATURE:

- ( A ) NAME/KEY: E1-CAT

## ( x i ) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Ala	Gly	Gly	Gly	Tyr	Trp	His	Thr	Ser	Gly	Arg	Glu	Ile	Leu	Asp	Ala	1	5	10	15
Asn	Asn	Val	Pro	Val	Arg	Ile	Ala	Gly	Ile	Asn	Trp	Phe	Gly	Phe	Glu	20	25	30	
Thr	Cys	Asn	Tyr	Val	Val	His	Gly	Leu	Trp	Ser	Arg	Asp	Tyr	Arg	Ser	35	40	45	
Met	Leu	Asp	Gln	Ile	Lys	Ser	Leu	Gly	Tyr	Asn	Thr	Ile	Arg	Leu	Pro	50	55	60	
Tyr	Ser	Asp	Asp	Ile	Leu	Lys	Pro	Gly	Thr	Met	Pro	Asn	Ser	Ile	Asn	65	70	75	80
Phe	Tyr	Gln	Met	Asn	Gln	Asp	Leu	Gln	Gly	Leu	Thr	Ser	Leu	Gln	Val	85	90	95	
Met	Asp	Lys	Ile	Val	Ala	Tyr	Ala	Gly	Gln	Ile	Gly	Leu	Arg	Ile	Ile	100	105	110	
Leu	Asp	Arg	His	Arg	Pro	Asp	Cys	Ser	Gly	Gln	Ser	Ala	Leu	Trp	Tyr	115	120	125	
Thr	Ser	Ser	Val	Ser	Glu	Ala	Thr	Trp	Ile	Ser	Asp	Leu	Gln	Ala	Leu	130	135	140	
Ala	Gln	Arg	Tyr	Lys	Gly	Asn	Pro	Thr	Val	Val	Gly	Phe	Asp	Leu	His	145	150	155	160
Asn	Glu	Pro	His	Asp	Pro	Ala	Cys	Trp	Gly	Cys	Gly	Asp	Pro	Ser	Ile	165	170	175	
Asp	Trp	Arg	Leu	Ala	Ala	Glu	Arg	Ala	Gly	Asn	Ala	Val	Leu	Ser	Val	180	185	190	
Asn	Pro	Asn	Leu	Leu	Ile	Phe	Val	Glu	Gly	Val	Gln	Ser	Tyr	Asn	Gly	195	200	205	
Asp	Ser	Tyr	Trp	Trp	Gly	Gly	Asn	Leu	Gln	Gly	Ala	Gly	Gln	Tyr	Pro	210	215	220	
Val	Val	Leu	Asn	Val	Pro	Asn	Arg	Leu	Val	Tyr	Ser	Ala	His	Asp	Tyr	225	230	235	240
Ala	Thr	Ser	Val	Tyr	Pro	Gln	Thr	Trp	Phe	Ser	Asp	Pro	Thr	Phe	Pro	245	250	255	
Asn	Asn	Met	Pro	Gly	Ile	Trp	Asn	Lys	Asn	Trp	Gly	Tyr	Leu	Phe	Asn	260	265	270	
Gln	Asn	Ile	Ala	Pro	Val	Trp	Leu	Gly	Glu	Phe	Gly	Thr	Thr	Leu	Gln	275	280	285	
Ser	Thr	Thr	Asp	Gln	Thr	Trp	Leu	Lys	Thr	Leu	Val	Gln	Tyr	Leu	Arg	290	295	300	
Pro	Thr	Ala	Gln	Tyr	Gly	Ala	Asp	Ser	Phe	Gln	Trp	Thr	Phe	Trp	Ser	305	310	315	320
Trp	Asn	Pro	Asp	Ser	Gly	Asp	Thr	Gly	Gly	Ile	Leu	Lys	Asp	Asp	Trp	325	330	335	
Gln	Thr	Val	Asp	Thr	Val	Lys	Asp	Gly	Tyr	Leu	Ala	Pro	Ile	Lys	Ser				

-continued

340

345

350

Ser Ile Phe Asp Pro Val  
355 358

## ( 2 ) INFORMATION FOR SEQ ID NO: 12:

- ( i ) SEQUENCE CHARACTERISTICS:  
( A ) LENGTH: 2293 base pairs  
( B ) TYPE: nucleic acid  
( C ) STRANDEDNESS: double  
( D ) TOPOLOGY: linear

( i i ) MOLECULE TYPE: DNA

- ( i x ) FEATURE:  
( A ) NAME/KEY: B1-CAT

( x i ) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

```
GGATCCACGT TGTACAAGGT CACCTGTCCG TCGTTCTGGT AGAGCGGCGG GATGGTCACC 60
CGCACGATCT CTCCTTTGTT GATGTCGACG GTCACGTGGT TACGGTTTGC CTCGGCCGCG 120
ATTTTCGCGC TCGGGCTTGC TCCGGCTGTC GGGTTCTGGT TGGCGTGGTG TCGGAGCAC 180
GCCGAGGCGA TCCCAATGAG GGCAAGGGCA AGAGCGGAGC CGATGGCACG TCGGGTGGCC 240
GATGGGGTAC GCCGATGGGG CGTGGCGTCC CCGCCGCGGA CAGAACCGGA TCGGAATAG 300
GTCACGGTGC GACATGTTGC CGTACCGCGG ACCCGGATGA CAAGGGTGGG TCGCGGGGTC 360
GCCTGTGAGC TGCCGGCTGG CGTCTGGATC ATGGGAACGA TCCCACCATT CCCCAGCAATC 420
GACGCGATCG GGAAGCAGGG GCGCGAGGCC GGACCGTGTG GTCGAGCCGG ACGATTGCGC 480
CATACGGTGC TGCAATGCCC AGCGCCATGT TGTCAATCCG CCAAATGCAG CAATGCACAC 540
ATGGACAGGG ATTGTGACTC TGAGTAATGA TTGGATTGCC TTCTTGCCGC CTACGCGTTA 600
CGCAGAATAG GCGACTGTAT GCGGTAGGTT GCGCTCCA GCGTGGGCTG GACATGCCTG 660
CTGCGAACTC TTGACACGTC TGGTTGAACG CGCAATACTC CCAACACCGA TGGGATCGTT 720
CCCATAAGTT TCGTCTCAC AACAGAATCG GTGCGCCCTC ATGATCAACG TGAAAGGAGT 780
ACGGGGGAGA ACAGACGGGG GAGAAACCAA CGGGGGATTG GCGGTGCCGC GCGCATTGCG 840
GCGAGTGCCCT GGTCTGCGGG TGATGCTGCG GGTCTGCGTC GTCGTGCGGG TGCTGGCATT 900
GGTTGCCGCA CTCGCCAACC TAGCCGTGCC GCGGCCGGCT CCGCGCGCGG GCGGCGGCTA 960
TTGGCACACG AGCGGCCGGG AGATCCTGGA CGCGAACAAC GTGCCGGTAC GGATCGCCGG 1020
CATCAACTGG TTTGGGTTTG AAACCTGCAA TTACGTCGTG CACGGTCTCT GGTACGCGGA 1080
CTACCGCAGC ATGCTCGACC AGATAAAGTC GTCGGGCTAC AACACAATCC GGCTGCCGTA 1140
CTCTGACGAC ATTCTCAAGC CGGGCACCAT GCCGAACAGC ATCAATTTTT ACCAGATGAA 1200
TCAGGACCTG CAGGGTCTGA CGTCCTTGCA GGTCAATGGAC AAAATCGTCG CGTACGCCGG 1260
TCAATATCGG CTGCGCATCA TTCTTGACCG CCACCGACCG GATTGCAGCG GGCAATCGGC 1320
GCTGTGGTAC ACGAGCAAGC TCTCGGAGGC TACGTGGATT TCCGACCTGC AAGCGCTGGC 1380
GCAGCGCTAC AAGGGAAACC CGACGGTCGT CGGCTTTGAC TTGCACAACG AGCCGCATGA 1440
CCCGGCCCTG TGGGGCTGCG GCGATCCGAG CATCGACTGG CGATTGGCCG CCGAGCGGGC 1500
CGGAAACGCC GTGCTCTCGG TGAATCCGAA CCTGCTCATT TTCGTGGAAG GTGTGCAGAG 1560
CTACAACGGA GACTCCTACT GGTGGGGCGG CAACCTGCAA GGAAGCCGGC AGTACCCGGT 1620
CGTGCTGAAC GTGCCGAACC GCCTGGTGTA CTCGGCGCAC GACTACGCGA CGAGCGTCTA 1680
CCCGCAGACG TGGTTCAGCG ATCCGACCTT CCCCACAAC ATGCCCGGCA TCTGGAACAA 1740
GAACTGGGGA TACCTCTTCA ATCAGAACAT TGCACCGGTA TGGCTGGGCG AATTCCGTAC 1800
```

-continued

GACACTGCAA	TCCACGACCG	ACCAGACGTG	GCTGAAGACG	CTCGTCCAGT	ACCTACGGCC	1860
GACCGCGCAA	TACGGTGCGG	ACAGCTTCCA	GTGGACCTTC	TGGTCCTGGA	ACCCCGATTG	1920
CGGCGACACA	GGAGGAATTC	TCAAGGATGA	CTGGCAGACG	GTCGACACAG	TAAAAGACGG	1980
CTATCTCGCG	CCGATCAAGT	CGTCGATTTT	CGATCCTGTC	TAATGATCTG	CATCGCCTAG	2040
CAGTCAACCG	TCCCCGTCGG	TGTCGCCGTC	TCCGTCGCCG	AGCCCGTCGG	CGAGTCGGAC	2100
GCCGACGCCT	ACTCCGACGC	CGACAGCCAG	CCCGACGCCA	ACGCTGACCC	CTACTGCTAC	2160
GCCACGCCC	ACGGCAAGCC	CGACGCCGTC	ACCGACGGCA	GCCTCCGGAG	CCCGCTGCAC	2220
CGCGAGTTAC	CAGGTCAACA	GCGATTGGGG	CAATGGCTTC	ACGGTAACGG	TGGCCGTGAC	2280
AAATTCGGGA	TCC					2293

We claim:

1. A DNA having the nucleotide sequence of SEQ ID NO. 12.
2. The DNA according to claim 1, encoding the amino acid sequence of SEQ ID NO. 11.
3. A vector carrying the DNA according to claim 1 and a vector sequence encoding either an origin of replication or an integration site for a host genome.
4. The vector according to claim 3 further comprising DNA encoding a signal sequence operably linked thereto.
5. The vector according to claim 3 further comprising exogenous regulatory sequences capable of causing expression of said DNA in a suitable host; wherein a microorganism comprising *Pichia pastoris* contains the vector.
6. A DNA according to claim 1 encoding the catalytic domain of E1 endoglucanase.
7. A DNA according to claim 1 encoding the catalytic domain and the linker peptide of E1 endoglucanase.

8. The DNA according to claim 6 further comprising at least one domain from a cellulase gene other than E1 endoglucanase.

9. The DNA according to claim 7 wherein the DNA encodes a protein having an endoglucanase activity.

10. A microorganism containing the vector of claim 5.

11. A method for producing an endoglucanase having the amino acid sequence of claim 2, comprising enzymatic cleavage of the cellulose binding domain of full size E1 endoglucanase.

12. A method for producing an endoglucanase having the amino acid sequence of claim 2, comprising inserting at least one stop codon after the sequence encoding the catalytic domain of full size E1 endoglucanase so that the linker peptide and cellulose binding domain of E1 endoglucanase are not expressed in the expression product.

13. The DNA according to claim 2 wherein the DNA encodes a protein having an endoglucanase activity.

\* \* \* \* \*